

Papillomavirus Probe and a process for in vitro
diagnosis of papillomavirus infections.

5

The invention concerns papillomavirus DNAs and more particularly the probes derived from these papillomaviruses, as well as the processes which use them for the in vitro diagnosis of papillomavirus infections.

The expression "papillomavirus" covers a great number of viruses having in common being considered responsible for several forms of viral infection ranging from relatively benign warts of the skin or mucous membranes to hyperplasias susceptible to degenerating into intra-epithelial neoplasms or into various forms of skin cancer. To be noted also among the papillomavirus infections are particularly the epidermodysplasias verruciformis which will sometimes be referred to hereafter by the expression "EV".

A certain number of types of papillomavirus have already been described. In the context of the present patent application, several new types and sub-types of papillomavirus will be described which have been isolated from warts or disseminated macular lesions, likely to lead to the development of precocious skin cancers in a large proportion of affected patients.

Recent studies have revealed the importance of immune factors and the major role of human papillomaviruses (HPV), to these factors added are the role previously described in the literature of various genetic factors and actinic radiations in the pathogenesis of papillomavirus infections.

The invention is the result of observations regarding the relative behavior of a great number of

T. 0 9 9 7 5 5 7 " 4 6 6 4

newly isolated papillomaviruses whose essential genomic characteristics will be defined below.

The study of a small number of EV cases has already lead to the characterization of 6 types of HPV after molecular cloning of their genomes (KREMSDORF, D. et al. 1982, J. Virol. 43, 436-447, and KREMSDORF et al. 1983, J. Virol. 48, 340-351). These HPV have been divided into three groups as a function of the lack of cross-hybridization or very weak cross-hybridization between the genomes of the different groups. The first group includes the HPV3a and 10 which are associated with the plane warts observed in certain EV patients and in the general population ; DNA sequences related to those of HPV3a have been found in the cancer of an EV patient. The second group includes HPV5, 8 and 12, the genomes of HPV5 and 8 having been detected in the cancers of EV patients. With the exception of a kidney transplant recipient presenting an immuno-suppression, who turned out to be infected by HPV5, the virus of the two latter groups have been detected only among EV patients, most of them being infected by several viruses. It should be noted that among the 14 types of HPV currently mentioned in the literature (bibliographic references 1-5, 8, 9, 13, 14, 16, 18-20 indicated further on), four turn out to be specifically associated with EV which is a rare disease.

The labors which have led to this invention and which have permitted the isolation of a large number of new types and sub-types of papillomavirus, create the possibility of more highly refined in vitro diagnostic techniques. More particularly, the invention provides perfected techniques for papillomavirus identification, for example, of those obtained from lesions or biopsy sections, and allows for more precise diagnoses which may also result in better prognoses with regard to the

possible evolution of the lesions in question.

As a general rule, it should be noted that, despite being very different from each other, the papillomaviruses have sizes of the order of 7000-8000
 5 base pairs. In addition, their genomes may nevertheless present certain degrees of homology. In what follows, reference will be made to evaluations of the percentage of homology between the various types and sub-types of papillomavirus, these homology percentages result from
 10 hybridization assays performed under conditions referred to as "non-stringent" or "non-strict", or under hybridization conditions called "stringent" ou "strict".

Among the papillomaviruses may be distinguished several types of papillomavirus, these may be disting-
 15 uished by their percentages of homology as measured under strict or stringent conditions. Papillomaviruses which, under these conditions, present homology percentages of less than 50 %, belong to different types. It may be noted in this regard, that the homology percent-
 20 ages between viruses of different types may even fall to zero under said strict or stringent conditions. Under these same conditions, viruses showing homology percentages of more than 50 % are considered as belonging to the same type and form the different sub-types within
 25 this same type.

Hybridization assays under non-strict or non-stringent conditions implies the mutual placing into contact of DNAs derived from two viral isolates under the following conditions as described by HEILMAN, C.A.
 30 et al. 1980, J. Virol. 36, 395-407, and CROISSANT et al. 1982, C.R. Acad. Sc. Paris, 294, 81-586 (hetero-duplex molecules).

Hybridization assays under strict or stringent conditions imply the placing into mutual contact of DNAs
 35 derived from two viral isolates under the conditions

T.D.G.T.T. " 4 6 9 2 8 5 5 0

described by KREMSDORF, D. et al. (1982, J. Virol. 43, 436-447, and J. Virol. 48, 340-351) and DAVIS, R.W. et al. 1971, Methods Enzymol. 21, 413-428 (hetero-duplex molecules).

5 Schematically, it may be noted that those papillomaviruses belonging to one same type presenting hybridizable sequences having virtually identical nucleotide sequences over 80 to 100 % of the totality of their respective lengths, these homologous sequences may
10 be reduced to 60 % or less among papillomaviruses of different types. The degree of identity or of analogy of the sequences from papillomaviruses of different types which mutually hybridize under non-strict or non-stringent conditions must obviously be less than in the case
15 of papillomaviruses of the same type.

The study to which the inventors proceeded has shown both that the degree of genetic heterogeneousness between diverse types of papillomaviruses was greater than previously recognized and at the same time that the
20 different types were found often to be associated with forms or variants of infections presenting a certain degree of specificity.

The invention consequently concerns not only the DNAs susceptible to being isolated from different new
25 papillomaviruses which have been isolated and the probes which can be partially or entirely constituted of these DNAs, but also mixtures or "cocktails" of papillomavirus types likely to be most effectively used for the diagnosis of diverse categories of infection, and of the
30 levels of risk to the patient which accompanies the discovery of a given papillomavirus. The number of papillomavirus probes described in the present application, to which may be added, as the case may be, those constituted from the genomic DNAs of papillomaviruses
35 which have already been previously isolated, and their

associations in the specific mixtures lend greater precision to the diagnosis, notably a greater discrimination between the diverse categories of infections which may be imputed to the diverse types of papillomavirus or which may be susceptible to develop under the effect of these latter types and, within a given category of infections, to give a better prognosis of the degree of risk that these latter could be transformed to more serious disease. For example, the invention aims to provide the means permitting, in the case of infections manifesting as epidermodysplasias verruciformis, to better appreciate the degree of risk that they may evolve towards skin cancer.

In a general manner and in an attempt to simplify the following presentation, the whole genomes of the papillomaviruses will be designated by the abbreviation "HPV-DNA".

With the same aim of simplification, reference will be made to appended drawings on which figure 20 physical restriction maps of HPV-DNAs including some from previously known papillomaviruses.

These physical maps give the positions of the fractionation sites of the various restriction endonucleases. The origin of the maps is generally constituted of the site of a single cut. The distances from this point of origin are expressed as a percentage of the total length of the genome. One unit of the map represents 1 % of the length of the genome.

The invention first of all concerns most specifically each of the HPV-DNAs chosen from among the total of the DNAs having sizes ranging between 7000 and 8000 base pairs and are characterized by the restriction maps appearing in the drawings which concern more particularly HPV-DNAs obtained from papillomaviruses and which correspond to the designations HPV-2d, HPV-10b,

HPV-14a, HPV-14b, HPV-15, HPV-17a, HPV-17b, HPV-19, HPV-20, HPV-21, HPV-22, HPV-23, HPV-24, HPV-28, HPV-29, HPV-31, and HPV-32, HPV-IP2 and HPV-IP4.

It goes without saying that the invention
5 equally extends its effects to HPV-DNAs which may be considered as belonging to the same types as those just enumerated.

The physical maps corresponding to the HPV-DNAs of newly characterized viruses are indicated by a solid
10 black circle on the drawings.

The invention equally concerns fragments of the preceding HPV-DNAs or capable of hybridizing with them, notably under strict conditions. Likewise it concerns those recombinant DNAs containing all or part of any of
15 the HPV-DNAs indicated above, and more particularly recombinant DNAs containing fragments corresponding to genes E1, E6-E7, L1, and L2 respectively or again fragments containing sequences corresponding to the inter-gene regions of these said HPV-DNAs. Lastly it
20 concerns the probes which may be constituted from these respective HPV-DNAs, and the processes of in vitro diagnosis using said probes.


Preparations of viral DNA have been selectively extracted (LUTZNER, M.A., 1983, Lancet ii:422-424) from
25 the products of scrapings of benign lesions of six European EV patients and two South American EV patients. The HPV-DNAs were purified by equilibrium centrifugation through cesium chloride gradients and/or by sedimentation through sucrose gradients in the presence of
30 ethidium bromide, according to the operating procedures previously described (articles by KREMSDORF, D. et al. mentioned above, and ORTH, G. et al. 1980, Cold Spring Harbor Conf. on Cell Proliferation 7:259-282). The DNA preparations were treated with restriction endonucleases
35 and the digestion products separated by electrophoresis

on agarose gels (articles by KREMSDORF et al. already mentioned). In addition, from the HPVs 5, 8 and 12 (KREMSDORF et al. articles again) and HPV-2 (HEILMAN, C.A. et al. 1980, J. Virol. 36:395-407, and ORTH, G. et al. 1980, Cold Spring Harbor conf. on Cell Proliferation 7:259-282) found in the verruca warts of one of the patients, eleven strains different from those from previously characterized types, were identified furnishing the major models for the restriction enzyme cleavage of the DNAs. The new HPV types were given a number and the sub-types of a type were given the same number followed by a letter according to the chronological order of their identification (COGGIN, J.R. et al. Cancer Res. 39:545-546). The genomes of the 11 new HPVs were cloned into Escherichia coli K12, strain C600 (KREMSDORF et al. article (1983) previously mentioned). The DNAs were inserted in the form of molecules of unitary length with the exception of two fragments of DNA from HPV-24 produced by endonuclease BamHI. They were inserted either into plasmid pBR322 (SUTCLIFFE, J.G., 1978, Nucleic Acids Res. 5:2721-2728), using the unique cleavage sites from AvaI, from BamHI, and from HindIII, or into a recombinant plasmid having integrated the HindIII B fragment from the DNA of HPV-5 (KREMSDORF et al. (1982) previously mentioned), which contains a unique SacI site. More particularly, the HPVs 17b and 22 were inserted in the form of DNA molecules of unitary length after splitting with an enzyme (SacI) which splits only once the DNA of HPVs and the recombinant plasmid pBR322 containing the HindIII B fragment from the DNA of HPV-5. The DNA of HPV-14a was inserted into the plasmid pBR322 in the form of a DNA molecule of unitary length after incomplete digestion of the viral DNA preparation by HindIII, an enzyme which produces two fragments of 96.1 and 3.9 % of the length of the genome.

T09T.T. 252350

the BamHI A and B fragments of HPV-24 (having sizes corresponding to 83.1 and 16.9 % respectively of the genome length) were inserted separately into plasmid pBR322.

5 The isolated clones and the corresponding HPV sources are listed in the appended Table I.



090557 44001

TABLE 1. ORIGIN OF DIAS FROM HPV CLONES

Patient ^a	Nationality	Type of DNA from HPV			Cloning ^c enzyme	Other types of HPV found in the patients
		Source ^b	cloned			
1	Polish	warts : knees	14a		Hind III	5
2	French	warts : hands	15		Dam III	
3	Columbian	macules : trunk	14b		Dam III	
4	Italian	macules : breast	17a		Dam III	5
			17b		Sao I	5
5	Dutch	macules : back	22		Sao I	5, 8, 17a
		macules : breast	19		Dam III	
6	Columbian	warts : hands	24		Dam III	5, 8, 24
7	Polish	warts : knees	20		Ava I	2, 12, 17a, 20
8	Polish	macules : forearm	21		Dam III	5, 8, 20
			23		Dam III	

To identify the recombinant plasmids, the electrophoretic mobilities of the digestion products of the recombinant DNAs and the non-cloned HPV-DNAs were compared after treatment with a mixture of two restriction endonucleases including the endonuclease used for the insertion of the viral sequences into the plasmid. The number and the sizes of the isolated fragments indicated that in each case the entire viral genomes were integrated. A heterogeneity of the DNA sizes was observed when non-cloned HPV-DNAs or those excised from plasmidic sequences, were analyzed by agarose gel electrophoresis (data not shown). The DNAs from HPVs 14b, 19, 20 and 21 have sizes similar to those of HPVs 3a, 5, 8 and 12 (around 7700 nucleotide pairs) (articles by KREMSDORF (1982) and ORTH (1980) already mentioned).

The sensitivity of cloned viral genomes to 14 restriction endonucleases was analyzed and physical maps established (Figures 1 through 10). The restriction maps of certain HPV-DNAs are repeated in some of the figures for reasons which will be explained further on. Between 22 and 33 cleavage sites have been localized according to the methods previously described (9). No apparent analogy could be detected between the maps with the exception of those of HPVs 14a and 14b, on the one hand (Figures 4a and 4b), and those of HPVs 17a and 17b, on the other (Figure 5). Among the 21 and 31 sites localized respectively on the DNAs of the HPVs 14a and 14b, fifteen turned out to be in common when one of the two BamHI cleavage sites of the DNA of HPV-14a was aligned with the unique BamHI cleavage site of the DNA of HPV-14b. In a similar manner, 21 of the 29 cleavage sites on the DNA of HPV-17a were equally found on the DNA of HPV-17b (out of 26 sites), when the sites of the unique BamHI cleavages were aligned.

No apparent analogy has been detected between

these maps and those previously established for the HPVs associated with EV (HPVs 3a, 5, 8, 9, 10 and 12) (8, 9, 16, 18 and 20), with warts on the skin (HPVs 1, 2, 4 and 7), with lesions of the muco-cutaneous or mucous membranes (HPVs 6b, 11a, 13 and 16), with the exception of the map for HPV-14a which is closely related to the map of an HPV isolated from a Japanese male with EV (24). This latter isolate differs from HPV-14a by one additional BamHI site and one additional HindII site, while the locations of the sites : AvaI, BamHI, BglI, EcoRI, HindII and HindIII were similar for the two viruses. Cross-hybridization experiments confirmed that these two viruses are indeed closely related.

It should be noted that some sites (those indicated by arrows) were not located. Cleavage sites differing by less than 2 % of the length of the genome were considered as retained (*). Enzymes which produced no splitting were : PvuI, SalI, and SmaI for the DNA of HPVs 14a and 23 ; PvuI, SacI, SalI and SmaI for the DNA of HPV-14b ; BglI, PvuI, SalI and SmaI for the DNA of HPVs 15, 17a and 17b ; BglI, SacI, SalI and SmaI for the DNA of HPV-19 ; EcoRI, PvuI, SacI and SmaI for the DNA of HPV-20 ; SacI and SmaI for the DNA of HPV-21 ; BamHI, BglI, PvuI, PvuII and SalI for the DNA of HPV-22 ; BglI, EcoRI, PvuI, SacI and SmaI for the DNA of HPV-24.

The existence of sequence homologies between the DNAs of the newly characterized HPV-DNAs as well as between these and the previously characterized HPV-DNAs of EV (HPVs 3a, 5, 8, 9, 10 and 12), of HPVs associated with skin warts (HPVs 1, 2, 4 and 7), and of HPVs associated with mucous membrane lesions (HPVs 6b, 11a, 13 and 16) has been studied. Hybridization experiments by attachment onto filter paper and DNA-DNA hybridization in saturated liquid phase followed by S1 nuclease digestion were carried out under the strict or stringent

20 unlabeled HPV-DNA.

25

30

35

0 0.2 0.4 0.6 0.8 1.0

cpm

unlabeled HPV-DNA

Total

Free

TABLE 2. DEGREE OF CROSS-HYBRIDIZATION BETWEEN DNAs FROM HPV, DETERMINED BY LIQUID PHASE HYBRIDIZATION.

Unlabeled		% of hybridization with DNA from HPV labeled with ³² P														
HPV DNA		3a	5	14a	14b	19	20	21	22	23	9	15	17a	17b	24	
1a		0.1	0.3	0.2	0.3	0	0.8	2.9	0	0	1.0	0.4	0.2	0	0	
11a		1.6	1.0	1.3	0	0.3	0.1	0.7	3.7	0	0.1	0.1	0.6	3.3	0	
3a		100	1.8	1.0	0	1.5	0.1	1.5	0	1.9	0.1	1.7	1.2	1.8	3.0	
10		32.3	ND	ND	0.1	0	0.1	1.9	3.0	ND	0	1.6	0.1	2.0	ND	
5		0.2	100	12.1	12.4	5.8	9.3	9.4	10.1	5.0	4.3	0.7	3.6	0	2.4	
8		1.1	15.7	9.9	13.4	8.5	5.6	5.0	7.1	5.0	3.5	1.5	3.2	3.8	0.1	
12		0.1	19.3	9.2	12.5	5.3	8.6	9.3	11.7	4.0	3.6	1.2	0.1	1.9	0	
14a		0.2	13.2	100	100.0	14.6	12.4	32.9	10.1	24.6	3.0	2.2	2.4	4.1	ND	
14b		ND	10.5	94.1	100	9.1	20.4	35.4	9.5	20.2	0	0	0	0	0	
19		ND	7.2	21.4	20.6	100	7.6	8.8	15.5	27.7	0	0	0	2.2	1.0	
20		ND	9.9	28.8	37.9	6.2	100	25.4	13.7	14.1	0	0	2.1	0	3.6	
21		ND	10.5	38.7	40.5	6.4	37.5	100	9.8	18.6	0.1	0	2.5	0.3	0	
22		ND	7.2	7.4	ND	17.3	7.2	10.0	100	17.9	0	0	0.1	0.1	0	
23		ND	ND	ND	ND	ND	ND	ND	21.2	100	0	0.5	0	0	0.1	
9		0.4	3.1	0.5	1.2	0	2.0	1.0	0	0	100	5.5	6.3	5.4	0	
15		0.4	3.3	2.1	3.3	0	0.1	0.8	0	0	7.8	100	22.5	21.6	0	
17a		0.7	1.2	1.4	2.0	0	0.1	0.1	0.8	1.4	7.6	19.5	100	92.7	0	
17b		ND	ND	ND	1.4	0	0.3	3.4	ND	ND	ND	ND	86.3	100	ND	
24		ND	ND	0.1	2.6	ND	ND	ND	0.8	0	0.2	0	1.1	1.1	100	

ND = not determined.

The absence or near absence of cross-hybridization between the genomes of HPVs 1, 2, 4, 6b, 7 and 11, and the HPV-DNAs of 32 P labeled newly cloned EV, or between unlabeled HPV-DNAs of EV and probes specific to HPVs 13, 16 and 18 is to be noted. In a similar manner, almost no cross-hybridization has been detected between the DNAs of HPVs 14a, 14b, 15, 17a, 17b, 19, 20, 21, 22, 23 and 24, and the DNAs of HPVs 1a and 11a by saturation reassociation (Table 2). The DNAs of newly cloned HPVs showed little or no cross-hybridization or showed less than 50 % cross-hybridization between each other and with the genomes of other HPVs associated with EV (HPVs 3a, 5, 8, 9, 10 and 12) with the exception of HPVs 14a and 14b on the one hand, and HPVs 17a and 17b on the other, which showed strong cross-hybridization. These observations justify the classification of the new viruses into nine new types (HPVs 14, 15, 17, 19, 20, 21, 22, 23 and 24) plus two sub-types of types 14 (HPV-14a and 14b) and 17 (HPV-17a and 17b).

Likewise, the different HPVs have been classed into groups based on their homologies (or lack of them) of sequences under the strict conditions of molecular hybridization. These groups, designated by the letters A to H, are listed in Table 3 which is appended. This Table mentions the diseases which have been diagnosed among the carriers of these HPVs (separately or in combination) and their oncogenic potential.

30

35




TABLE 1
CLASSIFICATION OF HPVs WHICH ARE THE OBJECT OF THIS PATENT APPLICATION, AS A FUNCTION OF THE DEGREE OF
HOMOLOGY OF THEIR NUCLEOTIDE SEQUENCES AS DETERMINED BY MOLECULAR HYBRIDIZATION UNDER STRICT CONDITIONS

Group ¹⁾	HPV types ²⁾	Homologies within the group	Associated diseases	Oncogenic potential	Probe mixture
A	1		Myrcemias	very weak	1
B	2		Verruca warts	weak	1
C	3, 10, 28*, 29*, 14 to 38 *		Plane warts Intermediary warts Actinic keratoses Bowen's disease	moderate ; a related virus, associated with intra-epithelial neoplasms and evolving skin cancers	2
D	4		Verruca warts	very weak	1
E	5, 8, 12, 14*, 19*, 20*, 21*, 22*, 23*	4 to 18 *	Epidermodysplasia verruciformis Actinic keratoses Bowen's disease Skin cancers	HPVs 5, 8 and 14 associated with cancers from E.V.; an related virus, associated with intra-epithelial neoplasms and evolving skin cancers	3, 4, 7
F	9, 15*, 17*, 24*	6 to 23 *	Epidermodysplasia verruciformis		5
G			Epidermodysplasia verruciformis		6
H	13, 31*		Oral epithelial hyperplasia ; oral leucoplasias		8
I	32*		Bowen's disease	Intraepithelial neoplasm and skin cancers	7, 9

1) Genomes of HPV types belonging to different groups, generally present no detectable sequence homology under strict molecular hybridization conditions. Genomes of HPV types belonging to the same group show less than 50 % sequence homology.

2) The new HPV types are indicated by an asterisk.

The DNAs of HPVs 5, 8, 12, 14, 19, 20, 21, 22 and 23 show between them levels of cross-hybridization (group homologies) varying from 5 to 38 % and show no notable cross-hybridization (4 to 13 %) except with the DNAs of HPVs 5, 8 and 12. These viruses thus form a part of a group of HPVs of EV previously defined (9).

Likewise, the DNAs of HPVs 9, 15 and 17 which show between them cross-hybridization of around 20 % and cross-hybridization of around 6 % with the DNA of HPV-9, equally belong to a group of EV HPVs previously described (9). The HPVs of types 13 and 31 may be considered as belonging to one same group. And last, the HPVs of types 1, 2, 4, 24 and 32 which show almost no homology with the genomes of the other HPVs, are considered as forming the first members of other groups which are distinct from each other and from the previous groups.

The invention concerns still more particularly the DNA fragments derived from the above described HPV-DNAs, and especially those corresponding to genes E6-E7, E1, L2, L1 and to their inter-gene regions. The relative positions and lengths of these various fragments in relation to the sites taken as origins (Figures 1 to 9) are indicated in Table 4 appended.




TABLE 4. PUTATIVE LOCATION OF THE PRINCIPAL GENES AND OF THE INTERGENE REGIONS ON THE PHYSICAL MAPS OF THE HPV GENOMES.

Type of HPV	Co-ordinates of the 5' and 3' ends, corresponding to genes				Intergene region
	E6 - E7	E1	L2	L1	
1	44 - 34.5	35 - 11	95 - 75.5	76.5 - 56	56 - 44.5
3	18.5 - 9	9.5 - 85.5	69.5 - 50	51 - 30.5	30.5 - 19
5	6.5 - 97	97.5 - 73.5	57.5 - 30	39 - 18.5	18.5 - 7
8	63 - 53.5	54 - 30	14 - 94.5	95.5 - 75	75 - 63.5
9	42 - 32.5	33 - 9	93 - 73.5	74.5 - 54	54 - 42.5
10a	49 - 39.5	40 - 16	0 - 80.5	81.5 - 61	61 - 49.5
10b	93 - 83.5	84 - 60	44 - 24.5	25.5 - 5	5 - 93.5
12	23.5 - 14	14.5 - 90.5	74.5 - 55	56 - 35.5	35.5 - 24
14	8.5 - 99	99.5 - 75.5	59.5 - 40	41 - 20.5	20.5 - 9
15	39.5 - 30	30.5 - 6.5	90.5 - 71	72 - 51.5	51.5 - 40
17	46 - 36.5	37 - 13	97 - 77.5	78.5 - 58	58 - 46.5
24	24.5 - 15	15.5 - 91.5	75.5 - 56	57 - 36.5	36.5 - 25
28	47.5 - 30	38.5 - 14.5	98.5 - 79	80 - 59.5	59.5 - 48
29	89.5 - 80	80.5 - 56.5	40.5 - 21	22 - 1.5	1.5 - 90
31	89 - 78.5	80 - 53.5	33.5 - 15.5	17.5 - 96.5	96.5 - 92.5

The localization of genes on the genome of HPV-1 has been deduced from the nucleotide sequence of this genome (patent of O. DANOS, M. KATINKA and M. YANIV). The physical maps of the genomes of the HPVs 3, 5, 8, 9, 10a, 12, 14, 15, 17 and 24 have been aligned relative to the physical map and the gene map of HPV-1, and that of HPV-31 relative to the physical map and gene map of HPV-6b (E. SCHWARZ and al, EMBO J., 1983, 2, 2341-2348), after electron microscope analysis of the hetero-duplex molecules formed under strict conditions (T_m -29 degrees C) or less strict (T_m -40 degrees C) of hybridization. The physical maps of HPVs 10b, 28 and 29 were lined up against the physical maps of HPVs 3a and 10a after juxtaposition of the retained restriction enzyme sites.

The values of the co-ordinates shown in table 4 give the position, on the physical maps presented in Figures 1-9, the 5' and 3' ends of the genome segments homologous with the genes E6 and E7, E1, L2 and L1 and with the inter-gene region relative to the genome of HPV-1a, or, in the case of HPV-31, relative to the genome of HPV-6b.

The inter-gene region (which includes the elements of regulation) and the adjacent E6 and E7 genes (corresponding probably to the major genes of transformation expressed in tumors) show no sequence homology detectable by electron microscope analysis of the hetero-duplex molecules formed (under non-strict hybridization conditions) between the genomes of HPV types belonging to different groups, or formed (under strict hybridization conditions) between the genomes of most of the HPV types belonging to the same group. Gene E1 (implicated principally in the replication of viral DNA) and gene L1 (coding for a major protein of the viral capsid bearing the virion's principal antigenic determinants) show sequence homologies detectable by the

analysis of hetero-duplexes formed (under non-strict hybridization conditions) between the genomes of HPV types belonging to different groups, or (under strict conditions) between the genomes of HPVs belonging to the same group.

Probes prepared from recombinant plasmids which include the E1 and L1 regions could theoretically permit the detection of the greatest number of HPV types by molecular hybridization experiments under strict or non-strict conditions according to the case. Probes prepared from recombinant plasmids which include the inter-gene region and the genes E6 and E7 permit the specific detection of one HPV type or related HPV types.

The L2 region (coding for a minor constituent of the viral capsid) shows a variable degree of conservation of nucleotide sequences among the different HPV types.

In what follows, the conditions under which the viruses HPV-IP2 and HPV-IP4 have been isolated will be described more precisely, as will the conditions under which the HPV-DNAs from these viruses were obtained.

Molecular cloning and characterization of a new type of HPV associated with neoplasms and genital cancers (HPV-IP2).

A new type of HPV has been demonstrated in the DNA extracted from a cancer of the cervix, by hybridization under non-strict conditions with a radioactive probe specific to HPV type 16. No cross-hybridization was detectable when the hybridization was carried out under the strict conditions of hybridization. A study of the sensitivity of the DNA of this HPV to a number of restriction enzymes has shown that the enzyme BglII cut once the viral DNA. After digestion of the DNA extracted from the tumor, by the endonuclease BglII, the fraction containing the DNA molecules of 8kb (size of a genome of

papillomavirus) were purified by centrifugation through
 a sucrose gradient. The 8kb molecules were inserted, at
 the BglII site, into a vector constituted of the plasmid
 PL15.5 (which includes a single cut site by BglII and by
 5 BamHI) which was inserted by it's BamHI site into the
 DNA of the bacteriophage lambda L47.1. After encapsida-
 tion of the recombinant DNA and infection of the host
 bacteria Escherichia coli strain (LA101), the areas of
 lysis corresponding to recombinant phages were detected
 10 by hybridization of the replicated infected bacterial
 cultures, with DNA from radioactive HPV-16, under non-
 strict conditions. Several recombinant bacteriophages,
 containing the totality of the viral sequences, have
 been isolated : the cutting of the phagic DNA by the
 15 insertion enzyme BglII results in a fragment of 8kb
 hybridizing with the HPV-16 under non-strict condi-
 tions ; the cutting of the DNA of the recombinant phages
 and of the DNA of the tumor of origin by a mixture of
 the enzymes BglII and PstI leads to the same 5 frag-
 20 ments, the total molecular weight of which is equal to
 the size of a genome of the papillomaviruses. The DNA of
 the new HPV was excised from the DNA of the recombinant
 bacteriophages, purified by electro-elution, and re-
 cloned in the plasmid PL15.5. A restriction map of the
 25 viral DNA has been established based on the sensitivity
 of this DNA to 18 restriction endonucleases, permitting
 21 cut sites to be localized (Figure 9). The map thus
 established is different from the map of the genomes of
 HPVs identified up til now. The sequence homology bet-
 30 ween the DNA of the new HPV and the DNA of the HPVs
 identified up til now were analyzed by molecular
 hybridization experiments on culture replicas under
 strict conditions. The homology detected was always less
 than 5 %, the greatest homology was detected with the
 35 genome of HPV-16. The new virus characterized from a

098755 44504

cervical cancer therefore constitutes a new type of HPV, provisionally called HPV-IP2.

The analysis, by electron microscopy, of the hetero-duplex molecules formed under different conditions between the DNA of HPV-IP2 and the DNA of HPV-1 permitted the alignment of the physical maps of these two genomes and the definition of the theoretical positions of the different genes of the DNA of HPV-IP2.

PUTATIVE POSITIONS OF THE PRINCIPAL GENES AND OF THE INTER-GENE REGION OF THE HPV-IP2 ON THE MAP OF THIS GENOME.

Co-ordinates of the ends : <u>5'</u> and <u>3'</u>		
E6-E7	62	71.5
E1	71	95
15 E2	95.5	11.5
L2	11	30.5
L1	31.5	52.
Inter-gene region	52	63.5

The use of radio-active probes prepared from the DNA of purified HPV-IP2 has permitted the determination of the pathogenic power of these viruses. The DNA of HPV-IP2 was found in one case of Bowenoid papula of the external genital organs, out of 14 cases studied ; in two cases out of 51 of invasive cervical cancer studied ; and in 1 case of cervical intra-epithelial neoplasm out of 28 studied. Thus HPV-IP2 constitutes a genito-tropic type of HPV with oncogenic potential whose frequency is a little less than that of HPV-18, and much less than that of HPV-16. It is necessary to include it in any mixture of HPV-DNAs intended for the preparation of molecular probes for the diagnosis of, or screening for, the types of HPVs presenting a risk of the development of genital neoplasms and, in particular, of cervical cancer.

Molecular cloning and characterization of a new type of HPV associated with pre-cancerous lesions of the skin (HPV-IP4).

A new type of HPV has been found in the DNA
 5 extracted from the biopsy of an actinic keratosis, a
 pre-cancerous skin lesion, by molecular hybridization
 under strict conditions, with a mixture containing
 radioactive probes specific for HPV types 5, 8 and 14.
 No cross-hybridization was detected when the hybridiza-
 10 tion was carried out with probes specific to types 1, 2,
 3, 7, 10, 13, 16, 18, 28, IP-1 (previously named HPV-
 31), IP-2 and IP-3 (previously named HPV-32).

A study of the sensitivity of this HPV to
 several restriction enzymes has shown that the enzyme
 15 EcoRI cuts the viral DNA once. After digestion of the
 DNA extracted from the biopsy by the endonuclease EcoRI,
 the fraction which contains the molecules of DNA of 8kb
 (size of a genome of papillomavirus) was purified by
 centrifugation through a sucrose gradient). The 8kb
 20 molecules were inserted, by the EcoRI site, into the DNA
 of the bacteriophage lambda gt¹⁰ wes. lambda B. After
 encapsidation of the recombinant DNA and infection of
 the host bacteria (Escherichia coli, strain LA101), the
 areas of lysis corresponding to the recombinant phages
 25 were detected by hybridization of the replicated
 infected bacterial cultures, with a radio-active mixture
 of DNAs from the HPVs 8 and 14 under non-strict condi-
 tions. Several recombinant bacteriophages, containing
 the totality of the viral sequences, have been isolated:
 30 the cutting of the phagic DNA by the insertion enzyme
 EcoRI results in an 8kb fragment hybridizing with the
 probe specific to HPVs 5, 8 and 14 under non-strict
 conditions ; the cutting of the DNA of the recombinant
 phages and the DNA of the original lesion by a mixture
 35 of the enzymes EcoRI and PstI results in the same 6


09987557 "11501
 TOSTT" 7528660

fragments, the total molecular weight of which equals the size of a genome of the papillomaviruses. The DNA of the new HPV was excised from the DNA of a recombinant bacteriophage, purified by electro-elution, and re-
5 cloned into the plasmid pSP65. A restriction map of the viral DNA was established from the sensitivity of this DNA to 15 restriction endonucleases, which has permitted the localization of 23 cut sites (Figure 10). The map thus established is different from the map of the
10 genomes of HPVs identified up til now. The sequence homology between the DNA of the new HPV and the DNAs of the HPVs identified up to now has been analyzed by molecular hybridization experiments on cultures under strict conditions. A homology, less than 50 %, has been
15 detected between the DNA of the new HPV and the DNA of certain types of HPVs previously identified in the lesions of epidermoplasia verruciformis (HPVs 5, 8, 12, 14, 19, 20, 21 and 25), but no homology was detected with other HPV types. The new virus characterized from
20 an actinic keratosis thus constitutes a new type of HPV provisionally named HPV-IP4.

The use of a radioactive probe prepared from the DNA of purified HPV-IP4 has permitted the demonstration of HPV-IP4 in 42 % of the 17 patients studied having
25 epidermodysplasia verruciformis and in x out of y biopsies of actinic keratosis analyzed. Because of its great frequency among patients of EV, a disease characterized by the frequent development of skin cancers, and because of its association with a fraction
30 of the lesions of actinic keratosis considered as precursors of spinocellular cancers of the skin. HPV-IP4 constitutes a type of dermo-tropic HPV with oncogenic potential. It is necessary to incorporate it into any mixture of HPV-DNAs intended for the preparation of
35 molecular probes for the diagnosis or screening of HPV

types constituting a risk of the development of cancerous or pre-cancerous skin lesions.

The invention still more particularly concerns the mixtures or cocktails of different HPV-DNAs (or probes containing these HPV-DNAs or sequences from them), susceptible to being used in combination to create means for the global diagnosis of the different forms of papillomavirus infections, possibly for use in the prognosis of the possible evolution of the infection. The preferred mixtures conforming to the invention are identified in Table 5 appended.



09977-1150
TOSTT 292960

TABLE 5
CHARACTERISTICS OF THE HPV-DNA MIXTURES USEABLE FOR THE DETECTION OF PAPILLOMAVIRUS INFECTIONS

Designation of the mixtures	Constitution ¹⁾	Diagnosable diseases
1	1, 2d*, 4	Skin or mucous membrane warts (especially verruca and plantar warts). Differential diagnosis of epidermodysplasia verruciformis.
2	3, 10a, 10b*, 28*, 29*	Plane warts or intermediary skin or mucous membrane warts. Intra-epithelial neoplasms and skin cancers. Differential diagnosis of epidermodysplasia verruciformis.
3	5, 17a*, 24*	Epidermodysplasia verruciformis. Intra-epithelial neoplasms and skin cancers.
4	5, 8, 12, 14a*, 14b*, 19*, 20*, 21*, 22*, 23*	Epidermodysplasia verruciformis.
5	9, 15*, 17a*, 17b*	Epidermodysplasia verruciformis.
6	24*	Epidermodysplasia verruciformis.
7	5, 8, 14b*, 32*	Skin cancers from epidermodysplasia verruciformis.
8	13, 31*	Intra-epithelial neoplasias and skin cancers.
9	32*	Oral epithelial hyperplasia ; differential diagnosis of oral epithelial neoplasms. Intra-epithelial neoplasms and skin cancers.

1) The new types of HPV are indicated by an asterisk.

This Table also indicates the natures of the affections susceptible to being more particularly diagnosed by the use of the mixtures figuring on the left side of the table. It should be noted that the grouping of the restriction maps in the appended figures 1 to 9 conform to the groupings given in the column headed "Composition" in table 5. This is equally the reason why some of the probes are reproduced several times in the different figures of the appended drawings.

Each of these mixtures may again be defined as including at least one of the new probes according to the invention. In other words, the diagnostic compositions according to the invention may be defined as containing :

- 1) at least the DNA of HPV-2d,
 - 2) at least the DNA of one of the HPVs 10b, 28 and/or 29,
 - 3) at least the DNA of one of the HPVs 17 and/or 24,
 - 4) at least the DNA of one of the HPVs 14, 15, 17, 19, 20, 21, 22 and/or 23,
 - 5) at least the DNA of one of the HPVs 15 and/or 17,
 - 6) the DNA of HPV-24,
 - 7) the DNA of HPVs 14 and 32,
 - 8) the DNA of HPV-31,
 - 9) the DNA of HPV-32,
- it being understood that the DNAs of the nine groups are chosen in such a manner as to in all circumstances be different from each other.

In view of the great diversity of HPVs susceptible of being isolated from the different forms of warts or other skin or mucous membrane lesions, it is nevertheless preferable, for the diagnosis of each type of disease mentioned in the table, to use mixtures containing more than one or two HPV-DNAs, since it has been recognized that other HPV-DNAs may equally intervene in

5

10

35

In the following is described other mixtures or cocktails of different HPV-DNAs (or probes containing these HPV-DNAs or sequences of them), susceptible of being used in combination for the realization of global diagnoses of the different forms of papillomavirus infections, perhaps for the purposes of determining a prognosis of the possible evolution of the affliction.

The preferred mixtures conforming to the invention are identified in the aforesaid table 5.

The aforesaid table equally indicates the natures of the affections susceptible of being more particularly diagnosed by the use of the mixtures figuring on the left side of the table. It is to be recalled that the restriction maps of other HPV-DNAs identified in the preceding table are contained in the Figures 1 through 9.

It should be noted that HPV-IP2 may be considered as particularly representative of the probes useable for the detection of the risk of developing genital neoplasms and, in particular, cervical cancers.

The invention therefore concerns more particularly yet the "kits" for diagnosis including at least 10 groups figuring in the groups numbered 1 to 10 in the table under the heading "Designation of the mixtures".

In the preceding, what was primarily under consideration was the use, as probes, of whole, cloned HPV-DNAs. These, however, could be replaced by cloned fragments of these different DNAs, notably by the genes E1 or L1 and by the genes E6-E7.

The basic principal of in vitro detections of HPV-DNAs will naturally involve hybridizations operated under strict or less strict conditions. A procedural example follows, it being of course understood that the diagnostic assays described should in no way be considered as limiting the conditions of use of the probes or

mixtures of the probes according to the invention.

The object of examinations involving probes prepared from mixtures of DNAs from cloned HPVs is to reveal an HPV and identify the type of HPV in a biopsy, in cells obtained by scraping a lesion, or in biopsy sections fixed with Carnoy's mixture (ethanol, chloroform, acetic acid 6:3:1) and included in paraffin. The examination requires the prior extraction of the DNA from the samples according to methods the principal of which is known and involves the analysis of this DNA by molecular hybridization under strict or less strict conditions with the aid of radioactive probes (labeled with ^{32}P or ^{35}S) prepared from mixtures of HPV-DNAs. In general, each examination requires the use of several mixtures of probes.

Several hybridization methods may be used. For example, the spot hybridization method includes, after denaturation of the DNA, the deposition of an aliquot of the DNA onto film supports (nitrocellulose or Genescreenplus), the hybridization of each film under the usual conditions with a mixture of probes, and the detection of the radioactive hybrids by contact exposition of the hybridized film onto radiographic film. Another possibility is replicated culture hybridization which involves agarose gel electrophoresis separation of the DNA fragments resulting from treatment of the DNA by restriction enzymes, the transfer of the fragments after alkaline denaturation onto films (nitrocellulose or Genescreenplus) and their hybridization under usual conditions with different mixtures of probes. The formation of radioactive hybrids is detected again by contact exposition of the hybridization support films onto radiographic film.

The radioactive probes are constituted of HPV-DNAs labeled by the "nick translation" method, or by

RNAs prepared by transcription of viral DNAs inserted
 into a vector, for example of the type SP6. The use of
 radioactive probes offers the advantage of great
 sensitivity, but this does not exclude the use of
 5 non-radioactive probes, for example biotinylated and
 able to be recognized by antibodies either themselves
 labeled or themselves recognizable by antibodies bearing
 enzymatic, fluorescent or other labeling.

The choice of probes depends on the nature of
 10 the patient sample. Thus, for example, in the case of a
 patient suspected of having EV, mixtures 1, 2, 3, 4, 5,
 6 and 7 would be used. Mixtures 1 and 2 would permit
 differential diagnosis between EV and skin warts. Probe
 3, which includes the most frequently detected member of
 15 each of the three groups of HPVs associated with EV, and
 probe 7, containing the DNAs of the HPVs associated with
 cancers of EV, would permit diagnosis of the majority of
 EV cases and, in particular, the identification of
 patients infected with the types of HPVs most likely to
 20 develop into cancers. The use of mixtures 4, 5 and 6
 would distinguish which type or types of HPV are
 infecting the same patient.

The invention thus again concerns the sets or
 "kits" containing a number of the probes indicated
 25 above, notably :

- either the representatives of each of the 19
types and sub-types of HPV-DNAs indicated above ;
- or mixtures of probes, preferably the diverse
groups or mixtures of probes which have been defined
 30 above, these kits being intended for diagnostic
 examinations in vitro by hybridization between the
 viral preparations obtained from patients, and the
 diverse groups of mixtures.

As it goes without saying and as it results
 35 already anyway from the preceding, the invention is in

no way limited to those of its embodiments and applications specifically anticipated, but, on the contrary, embraces all variants, notably the reference in the claims to a designation HPV-DNA followed by a specific number, and to which corresponds an HPV-DNA whose restriction map has been provided in the drawings, is to be understood as signifying that these claims cover all HPV-DNAs which have in common with this particular HPV-DNA, the power to be classed in the same type, according to the definition which was given above, and all the more to HPV-DNAs belonging to the same sub-type.

It is also to be noted, particularly with regard to the DNA derived from HPV-32 which appears in the drawings, that it is not split by any of the following enzymes : AvaI, BalI, BamHI, ClaI, EcoRI, HindIII, NdeI, NruI, PvuII, SacI, SalI, SmaI, TthIII, or XmaI.

Note, too, that the following recombinant DNAs were filed with the C.N.C.M. (National Collection of Cultures of Micro-organisms of the Institut Pasteur in Paris) the 30th of November 1984, under the numbers listed here :

	pBR322/HPV2d.....	No. I-379
	pBR322/HPV10bA.....	No. I-380
	pBR322/HPV10bB.....	No. I-381
25	pBR322/HPV14a.....	No. I-382
	pBR322/HPV14b.....	No. I-383
	pBR322/HPV15.....	No. I-384
	pBR322/HPV17a.....	No. I-385
	pHPV5 HindIIIB/HPV17b.....	No. I-386
30	pBR322/HPV19.....	No. I-387
	pBR322/HPV20.....	No. I-388
	pBR322/HPV21.....	No. I-389
	pHPV5 HindIIIB/HPV22.....	No. I-390
	pBR322/HPV23.....	No. I-391
35	pBR322/HPV24a.....	No. I-392

	pBR322/HPV24b.....	No. I-393
	pBR322/HPV28.....	No. I-394
	pBR322/HPV29.....	No. I-395
	pBR322/HPV31.....	No. I-396
5	pSP64/HPV32.....	No. I-397
	pLI55/IP2.....	No. I-450
	pSP65/IP4.....	No. I-449

The invention yet more particularly the products of the expression of genes E6 and E7 from the different papillomaviruses which have been evoked in the preceding and which may be used as the active principal of vaccines able to induce, when administered in effective doses, the resistance of the host to the development of neoplasms associated with papillomaviruses.

15 The invention equally concerns serums susceptible to being obtained by immunization of a mammal, serums which may be used for the preparation of serums administrable in effective doses to a patient, notably parenterally, these serums then being able to provoke a remission of the infections induced by the corresponding types or sub-types of papillomaviruses.

It is pointless to emphasize the capacity of the specialist to obtain the products of the polypeptidic expression of the kind in question, notably by the techniques of genetic engineering consisting of the incorporation of the E6 and/or E7 sequences into a vector under the control of an appropriate promoter, then the transformation of a cellular host, of which the papillomaviruses are likely to recognize the promoters in question and to express the sequences associated with it.

The invention thus also concerns the compositions for pharmaceutical usage containing the principles of the kind in question (expression products or corresponding antibodies), in association with a

T.O.S.T. 454856

5

10



BIBLIOGRAPHY

- (1) Dürst, M. et al., 1983, Proc. Natl. Acad. Sci. U.S.A., 80:3812-3815.
- (2) Coggin, J.R., Jr. et al., 1979, Cancer Res., 39:545-546.
- (3) Gissmann, L. et al., 1982, J. Virol. 44:393-400.
- (4) Green, M. et al., 1982, Proc. Natl. Acad. Sci. U.S.A. 79:4437-4441.
- (5) Heilman, C.A. et al., 1980, Virol. 36:395-407.
- (6) Jablonska, S. et al., 1972, Cancer Res., 32:583-589.
- (7) Jablonska, S. et al., 1982, Springer Semin. Immunopathol. 5:33-62.
- (8) Kremsdorf, D. et al., 1982, J. Virol. 43:436-447.
- (9) Kremsdorf, D. et al., 1983, J. Virol. 48:340-351.
- (10) Lutzner, M.A. et al., 1978, Bull. Cancer, 65:169-182.
- (11) Lutzner, M.A. et al., 1983, Lancet ii:422-424.
- (12) Migozzi, M. et al., 1965, Bull. Soc. Franc. Derm. Syph. 72:747-748.
- (13) Orth, G. et al., 1980, Cold Spring Harbor Conf. Cell Proliferation, 7:259-282.
- (14) Orth, G. et al., 1981, J. Invest. Dermatol. 76:97-102.
- (15) Orth, G. et al., 1979, Cancer Res. 39:1074-1082.
- (16) Ostrow, R.S. et al, 1982, Proc. Natl. Acad. Sci. U.S.A., 79:1634-1638.
- (17) Ostrow, R.S. et al., 1983, Ann. Acad. Dermatol. 8:398-404.
- (18) Pfister, H. et al., 1983, Cancer Res. 43:1436-1441.
- (19) Pfister, H. et al., 1983, J. Virol. 47:363-366.
- (20) Pfister, H. et al., 1981, Int. J. Cancer, 27:645-650.
- (21) Rueda, L.A. et al., 1976, Med. Cut. I.L.A. 2:113-136.
- (22) Ruiter, M. et al. J. Invest. Dermatol., 47:247-252.
- (23) Sutcliffe, J.G., 1978, Nucleic Acids Res. 5:2721-2728.
- (24) Tsumori, T. et al., 1983, J. Gen. Virol. 64:967-969.